



Questionnaire survey on use of placebo

Uriel Nitzan and Pesach Lichtenberg

BMJ 2004;329:944-946; originally published online 17 Sep 2004;
doi:10.1136/bmj.38236.646678.55

Updated information and services can be found at:
<http://bmj.com/cgi/content/full/329/7472/944>

These include:

Data supplement

"English translation of the questionnaire"
<http://bmj.com/cgi/content/full/bmj.38236.646678.55/DC1>

References

This article cites 7 articles, 2 of which can be accessed free at:
<http://bmj.com/cgi/content/full/329/7472/944#BIBL>

5 online articles that cite this article can be accessed at:
<http://bmj.com/cgi/content/full/329/7472/944#otherarticles>

Rapid responses

8 rapid responses have been posted to this article, which you can access for free at:
<http://bmj.com/cgi/content/full/329/7472/944#responses>

You can respond to this article at:
<http://bmj.com/cgi/eletter-submit/329/7472/944>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top left of the article

Topic collections

Articles on similar topics can be found in the following collections

[Other Ethics](#) (1393 articles)
[Patient - caregiver communication \(including Patient education\)](#) (432 articles)
[Patient - caregiver relationships](#) (190 articles)

Notes

To order reprints follow the "Request Permissions" link in the navigation box

To subscribe to *BMJ* go to:
<http://resources.bmj.com/bmj/subscribers>

Questionnaire survey on use of placebo

Uriel Nitzan, Pesach Lichtenberg

Herzog Hospital
and Hadassah
School of Medicine,
Department of
Psychiatry, POB
35300, Jerusalem
91351, Israel
Uriel Nitzan
medical student
Pesach Lichtenberg
lecturer

Correspondence to:
P Lichtenberg
licht@cc.huji.ac.il

BMJ 2004;329:944-6

Abstract

Objectives To gauge the frequency and circumstances of use of placebo in clinical practice and the attitudes towards its use among those who administer it.

Design Retrospective questionnaire.

Setting Two large hospitals and various community clinics in the Jerusalem area.

Participants 31 physicians working in hospital inpatient and outpatient departments, 31 head nurses working in hospital inpatient departments, and 27 family physicians working in community clinics.

Main outcome measures Self report of frequency and circumstances of, and attitudes towards, use of placebo.

Results Among the 89 respondents, 53 (60%) used placebos (95% confidence interval 49% to 70%). Among users, 33 (62%) prescribed a placebo as often as once a month or more; 36 (68%) told patients they were receiving actual medication; 15 (28%) considered that placebos were a diagnostic tool; and 48/51 (94%) reported that they found placebos generally or occasionally effective.

Conclusion Most practitioners questioned in this study continue to use placebos. Used wisely, placebos might have a legitimate place in therapeutics. Wider recognition of the practice and debate about its implications are imperative.

Introduction

How common is the use of placebos in clinical practice? From the dearth of discussion in the medical literature—almost all references to placebo from a Medline search refer to a research context—one might surmise that the clinical use of placebos is exceedingly rare. The deception involved in administering a placebo certainly raises ethical questions, and some institutions have banned its use. Yet informal discussions with colleagues, as well as first hand observation of clinical activity in various medical departments, suggest the practice still occurs. The only other report that we could locate of placebo use in a clinical context goes back a quarter of a century.¹ The authors found that the placebo was rarely (about once a year per physician) and inappropriately used.

We reappraised how frequently and in what circumstances physicians and hospital nurses use placebos in a clinical setting, how they understand the mechanisms of actions, their views on ethics, and whether they find placebos useful.

Methods

Questionnaire

We developed a questionnaire on attitudes and experience with placebos. We sought information on basic demographics, the frequency of placebo use, the circumstances of its administration, accompanying beliefs about mechanisms of action, and ethical positions. The questionnaire was administered in Hebrew (see bmj.com for an English translation).

Recruitment

We approached three groups of physicians and nurses, who might be expected to differ in their attitudes towards the use of a placebo and its legitimacy as a therapeutic tool. Of around 110 physicians and nurses approached, 89 agreed to participate in the survey, all of whom received and returned completed questionnaires.

Senior physicians working in hospital inpatient and outpatient departments in medical and surgical specialties and subspecialties—The bulk of the work of these respondents was with inpatients, with one or two half days a week spent in the associated hospital based outpatient clinic. So as not to collect duplicate data for the same inpatient department, we included only one physician from each department. We covered all 31 inpatient departments at two major hospitals in Jerusalem and therefore included 31 respondents in this category. Ten physicians declined to receive the questionnaire.

Head nurses working in the same hospital inpatient departments as senior physicians—Here too we had 31 respondents, one per department. As the nurses are responsible for dispensing medications on inpatient wards, we included them to get a picture of what actually takes place in the hospital services. All nurses approached agreed to participate in the survey.

Family physicians working in community clinics—These respondents were recruited from a weekly gathering of about 40 family physicians working in Jerusalem. Of these, 27 agreed to receive the questionnaire and all completed it. As we expected physicians working in the same clinic to display independent prescribing habits, we allowed more than one physician from a single clinic to respond.

Statistical methods

To assess the precision of the estimated proportion of placebo use in clinical practice, we calculated 95% confidence intervals. We used Pearson χ^2 test and Fisher's exact test to test the significance of the association between two qualitative parameters. $P \leq 5$ was considered significant.

Results

Table 1 shows the composition of the group. We have summarised the data from the questionnaire under six headings.

Frequency—When we planned the study we assumed that the use of placebo was not widespread and would not exceed 10%. Among our 89 respondents, however, 53 (60%) admitted using a placebo (95% confidence interval 49% to 70%). The age and sex of respondent did not affect results. In total, 53% of doctors and 71% of nurses reported using a placebo. Among users, 33 (62%; 37% of the total sample) used a placebo as often as once



An English translation of the questionnaire can be found on bmj.com

This article was posted on bmj.com on 17 September 2004: <http://bmj.com/cgi/doi/10.1136/bmj.38236.646678.55>

Table 1 Demographic and professional data for respondents to questionnaire on use of placebos

	No (%) [*]
Mean (SD) age (years)	39.4 (9.3)
Men	37 (42)
Women	52 (58)
Profession:	
Doctor	58 (65)
Nurse	31 (35)
Mean (SD) professional experience (years)	13.0 (9.7)
Primary work setting:	
Internal medicine department	32 (36)
Surgical department	30 (34)
Family medicine clinic	27 (30)

^{*}Unless stated otherwise.

a month or more. Differences between physicians and nurses in reported use and frequency of use of placebo did not attain significance.

Perceived therapeutic value—Of those who used a placebo, most (48 of 51 who answered the question, or 94%) found that it was either generally (17, or 33%) or occasionally (31, or 61%) effective.

Information given to patients—Of those using the placebo, 36 (68%) tell the patient that he or she is receiving a real medicine, and nine (17%) say nothing at all. The rest either identify the placebo as such (two, or 4%) or tell the patient that he or she is receiving a non-specific medicine (six, or 11%).

Circumstances of use—We found a wide range of applications for placebo (table 2). Placebos were given in the form of saline infusions or intramuscular injections; paracetamol or vitamin C tablets instead of the ordinarily prescribed medication; sugar or artificial sweetener pills; or prepared placebo tablets. The medical conditions for which the placebos were used included anxiety, pain (including abdominal), agitation, vertigo, sleep problems, asthma, contractions in labour, withdrawal from recreational drugs, and angina pectoris (when the blood pressure was too low to allow for vasodilators). The stated value as a diagnostic tool, referred to in table 2, was to distinguish organic from psychogenic or simulated arthralgia, seizure disorder, and abdominal or other pain.

Ethical stance—Of 79 responses on ethics, only four (5%) thought that the use of placebos should be categorically prohibited. Most of the others considered placebo use conditional on certain circumstances, such as prior experience (26, or 33%), notifying patients of receipt of a placebo (23, or 29%), or evidence from research that the placebo was effective (19, or 24%).

Perceived mechanism of action—Respondents were permitted to propose more than one mechanism of

action for placebos. Of 83 responses, most (62, or 75%) attributed the effect purely to psychological mechanisms. An additional nine (11%) respondents suggested a combination of psychological and biochemical effects.

Discussion

Principal findings

We attempted to gauge the extent of placebo use in clinical work. Despite general disapproval in the medical literature,^{2,3} such use continues among 60% of our respondents. The circumstances varied but included a wide variety of clinical situations. Indeed, as only one in 20 would prohibit the placebo in all circumstances the potential for placebo prescriptions is even greater than the actual extent of use. We also found that most practitioners who use placebos claim effectiveness for the treatment in some or most cases. In light of this finding, it seems likely that the many such practitioners will continue to prescribe placebos.

Many physicians relate to the placebo as a diagnostic tool. This indicates a persistence of long discredited notions of a separation between mind and body. A placebo can assuage pain. Even in a meta-analysis that raised questions about the actual existence of a placebo effect the authors concurred that placebos can have analgesic potency.⁴ The physician who nevertheless uses a placebo diagnostically is at risk of reaching unfounded conclusions, to the detriment of his or her patients.

Strengths and weaknesses of study

Because we investigated all medical and surgical inpatient departments at two hospitals and chose only one senior physician and nurse from each, our findings are quite comprehensive. A weakness of our paper, however, is that we rely on self reports made retrospectively. This can be a problem when respondents are asked to look back and estimate the frequency of a particular behaviour. The fact that a particular respondent uses a placebo in a clinical context is not likely to be misremembered, however, even if the frequency is misjudged. Moreover, in light of the suspect moral validity of such treatment, we would anticipate that placebo use would be understated in the responses to the questionnaire. If so, our finding that placebo prescribing is a widespread practice cannot be doubted.

Previous study

In a previous study, the extent of placebo use was found to be considerably less, about one prescription a year per physician.¹ In our study over a third of the respondents reported using a placebo once a month or

Table 2 Circumstances in which placebo was administered. Figures are number (percentage of those who reported use) of respondents^{*}

	Hospital based physicians	Nurses	Family physicians	Total
After "unjustified" demand for medication	8 (42)	8 (36)	7 (58)	23 (43)
To calm patient	4 (21)	9 (41)	7 (58)	20 (38)
As analgesic	4 (21)	11 (50)	5 (42)	20 (38)
As diagnostic tool	6 (32)	4 (18)	5 (42)	15 (28)
As adjunctive therapy	3 (16)	3 (14)	6 (50)	12 (23)
For non-specific complaints	0	2 (9)	7 (58)	9 (17)
To buy time before next regular dosage of medication	2 (11)	5 (23)	1 (8)	8 (15)
To get patient to stop complaining	1 (5)	2 (9)	3 (25)	6 (11)

^{*}53 respondents reported that they use placebo (19 hospital based physicians, 22 nurses, and 12 family physicians). Each respondent was permitted to cite more than one circumstance in which he or she used placebo.

more, and nearly 60% at least once a year. Though the differing methods of the data collection render comparisons across these studies difficult, our findings suggest that the use of the placebo is increasing. The former study recorded actual prescriptions and was limited to actual placebo pills or saline, while we studied the self reports of healthcare providers, and, in certain circumstances, paracetamol and vitamin pills were also counted as placebos.

Implications

Some have advocated banning the clinical use of placebos because of the deception involved in administration and the possible harm to the doctor-patient relationship.^{2,3} Others have suggested guidelines for the proper use of placebos without violating the patient's trust and autonomy.⁵⁻⁷ Our study shows that administration of placebos for clinical purposes continues. Clearly, wider recognition of the practice, and debate about its implications, are needed. Further investigations into the extent and nature of use should be conducted, particularly in a clinical context where the placebo's effect may differ from that found in randomised controlled trials.^{8,9} Moreover, though Israeli medicine is taught and practised as elsewhere in the Western world, similar surveys in other geographical areas may reveal cross cultural differences.

Contributors: UN had the original idea for the study and compiled the data. Both authors jointly developed the questionnaire. PL wrote most of the article and is guarantor.

Funding: None.

Competing interests: None declared.

Ethical approval: Not required.

What is already known on this topic

No study has recently attempted to assess the use of placebos in clinical settings

Placebos may be effective in some areas, but their use raises ethical issues

What this study adds

Three in five clinicians continue to use placebos

Clinicians believe that some patients benefit from placebo treatment

The role of placebo treatment, its mechanisms, and its ethics need to be the subject of wider medical education and debate

- 1 Goodwin JS, Goodwin JM, Vogel AV. Knowledge and use of placebos by house officers and nurses. *Ann Intern Med* 1979;91:106-10.
- 2 De Deyn PP, d'Hooge R. Placebos in clinical practice and research. *J Med Ethics* 1996;22:140-6.
- 3 Hill J. Placebos in clinical care: for whose pleasure? *Lancet* 2003;362:254.
- 4 Hrobjartsson A, Gotzsche PC. Is the placebo powerless? An analysis of clinical trials comparing placebo with no treatment. *N Engl J Med* 2001;344:1594-602.
- 5 Brown WA. Placebo as a treatment for depression. *Neuropsychopharmacology* 1994;10:265-9.
- 6 Lione A. Ethics of placebo use in clinical care. *Lancet* 2003;362:999.
- 7 Lichtenberg P, Heresco-Levy U, Nitzan U. The ethics of the placebo in clinical practice. *J Med Ethics* (in press).
- 8 Vase L, Riley JL, Price DD. A comparison of placebo effects in clinical analgesic trials versus studies of placebo analgesia. *Pain* 2002; 99:443-52.
- 9 Hyland ME. Using the placebo response in clinical practice. *Clin Med* 2003;3:347-50.

(Accepted 13 August 2004)

doi 10.1136/bmj.38236.646678.55

Are written responses to some referrals to a general haematology clinic acceptable?

Allison Tso, Lucy Harris, Tim Littlewood

Department of Haematology, John Radcliffe Hospital, Oxford OX3 9DU

Allison Tso
senior house officer in medicine

Lucy Harris
medical student

Tim Littlewood
consultant haematologist

Correspondence to:
T Littlewood
tim.littlewood@orh.nhs.uk

BMJ 2004;329:946-7

Each year about 1000 patients are referred to the department of haematology in Oxford. Excluded from this number are referrals for problems related to coagulation, which are seen at the Oxford Haemophilia Centre. We wondered whether a written reply to the general practitioner would provide the same quality of healthcare advice more quickly, and with less inconvenience to the patients, than seeing patients in a hospital outpatient clinic.

Methods and results

One consultant (TL) received 274 letters of referral between 1 November 2001 and 1 January 2003 and wrote a response to 121 (table). We subsequently sent each of the general practitioners who had received a written response a questionnaire (box) with a copy of their original referral letter and a copy of the written reply.

TL sent written responses if the patient neither required further investigation (such as a bone marrow biopsy) nor treatment that would better be done in the haematology department and if no evidence indicated a serious underlying illness, such as malignancy. For

Questionnaire

Was a written reply offering advice rather than an outpatient appointment acceptable to you? Yes/No
Was the advice given helpful to you? Yes/No
Would you be satisfied with a written response in the future if thought appropriate? Yes/No
Do you know if the patient was satisfied with a written response? Yes/No
Invited other comments

example, a mild macrocytosis (mean cellular volume less than 105 fl) without accompanying cytopenia was the commonest referral for which TL sent a written response. In all patients, the blood film had been examined (and was normal apart from the mild macrocytosis) and advice was offered about checking for possible causes of macrocytosis (including vitamin B-12 or folate deficiency, liver disease, hypothyroidism, drug related causes, and excess alcohol consumption), if these factors

This article was posted on bmj.com on 1 October 2004: <http://bmj.com/cgi/doi/10.1136/bmj.38253.703553.F7>